

## ORAL PRESENTATION

## Open Access

# Sagittal and pelvic parameters analysis in patients with adolescent idiopathic scoliosis

S Donzelli\*, F Zaina, S Negrini

From 8th International Conference on Conservative Management of Spinal Deformities and SOSORT 2011 Annual Meeting  
Barcelona, Spain. 19-21 May 2011

## Background and purpose

The sagittal alignment of the spine and pelvis in adolescent idiopathic scoliosis is poorly defined in the literature [1-8]. The purpose of this study was to assess the sagittal alignment in scoliosis patients according to curve degree and type.

## Material and methods

Sagittal parameters of the spine and pelvis were analysed in lateral standing radiographs of 256 adolescents (13.7±5 years, curve range 4-57°) and compared with statistically normal values (NV) in adolescents found in the literature: thoracic kyphosis TK (NV 22-66°), lumbar lordosis LL (NV 24-72°), pelvic incidence PI (NV 27-71°), sacral slope SS (NV 25-57°) and pelvic tilt PT (NV -8-16°). Lateral standing radiographs were matched with anteroposterior radiograph. Patients were classified according to the entity of scoliosis curves, age, gender and risser score.

## Results

There is a weak negative correlation (0.2) between scoliosis and kyphosis. Over 20° Cobb PI increased, mainly due to an increase of the SS. In our population we had low PI and SS, but mainly in less than 20° curves than in higher scoliosis; on the contrary, PT was high in all children. Analysing curves type and decrease of SS we found that this occurs more frequently in patients with double curves (thoracic and thoracolumbar).

## Conclusions

PI increases through life, and curves degree worsen with growth, and this influence our results. Patients with spinal deformities have a positive sagittal balance and signs of pelvic retroversion such as decreased SS. According to our

data this situation occurs to patients with thoracic and thoracolumbar curves.

Published: 27 January 2012

## References

1. Cil Akin, Yazici Muharrem, Uzunucugil Akin, Kandemir Utku, Alanay Ahmet, Alanay Yasemin, Emre Acaroglu R, Surat Adil: **The evolution of sagittal segmental alignment of the spine during childhood.** *Spine* 2004, **30**(1):93-100.
2. Mac-Thiong JM, Labelle H, Berthodnaud E, Betz RR, Roussouly P: **Sagittal spinopelvic balance in normal children and adolescents.** *Eur Spine J* 2007, **16**:227-234.
3. Vaz G, Roussouly P, Berthodnaud E, Dimnet J: **Sagittal morphology and equilibrium of pelvis and spine.** *Eur Spine J* 2002, **11**:80-87.
4. Roussouly P, Gollogly S, Berthodnaud E, Dimnet J: **Classification of the normal variation in the sagittal alignment of the human lumbar spine and pelvis in the standing position.** *Spine* 2005, **30**(3):346-353.
5. Roussouly P, Gollogly S, Nosedà O, Berthodnaud E, Dimnet J: **The vertical projection of the sum of the ground reactive forces of a standing patient is not the same as the C7 plumb line: a radiographic study of the Sagittal Alignment of 153 asymptomatic volunteers.** *Spine* 2006, **31**(11):E320-E325.
6. Mac-Thiong JM, Roussouly P, Berthodnaud E, Guigui P: **Sagittal parameters of global spinal balance normative values from a prospective cohort of seven hundred nine caucasian asymptomatic adults.** *Spine* 2010, **35**(22):E1193-E1198.
7. Mendoza-Lattes S, Ries Z, Gao Y, Weinstein SL: **Natural history of spinopelvic alignment differs from symptomatic deformity of the spine.** *Spine* 2010, **35**(16):E792-E798.
8. Schwab FJ, Smith V, Biserni M, Gamez L, Farcy JPG, Pagala M: **Adult Scoliosis: a quantitative radiographic and clinical analysis.** *Spine* 2002, **27**(4):387-392.

doi:10.1186/1748-7161-7-S1-O15

**Cite this article as:** Donzelli et al.: Sagittal and pelvic parameters analysis in patients with adolescent idiopathic scoliosis. *Scoliosis* 2012 **7** (Suppl 1):O15.